



## SHANK3 gene

SH3 and multiple ankyrin repeat domains 3

### Normal Function

The *SHANK3* gene provides instructions for making a protein that is found in many of the body's tissues but is most abundant in the brain. The SHANK3 protein plays a role in the functioning of synapses, which are the connections between nerve cells (neurons) where cell-to-cell communication occurs. Within synapses, the SHANK3 protein acts as a scaffold that connects neurons, ensuring that the signals sent by one neuron are received by another.

The SHANK3 protein is also involved in the formation and maturation of dendritic spines. Dendrites are specialized extensions from neurons that are essential for the transmission of nerve impulses. Dendritic spines are small outgrowths from dendrites that further help transmit nerve impulses and increase communication between neurons.

### Health Conditions Related to Genetic Changes

#### 22q13.3 deletion syndrome

The characteristic signs and symptoms of 22q13.3 deletion syndrome, which is also commonly known as Phelan-McDermid syndrome, are caused by a deletion near the end of the long (q) arm of chromosome 22. The chromosomal region that is typically deleted is thought to contain many genes, including the *SHANK3* gene. As a result of the deletion, people with this condition have only one copy of the *SHANK3* gene in each cell instead of the usual two copies.

Researchers believe that a deletion of the *SHANK3* gene and a reduction in the amount of SHANK3 protein produced is responsible for many of the features of 22q13.3 deletion syndrome. A decrease in the functioning of synapses and cell-to-cell communication between neurons caused by a lack of SHANK3 protein is thought to contribute to the developmental delay, intellectual disability, and absent or severely delayed speech characteristic of people with 22q13.3 deletion syndrome.

#### other disorders


At least seven *SHANK3* gene mutations have been found in people who have the features of autism or similar conditions known as autism spectrum disorders. These disorders are characterized by impaired communication and socialization skills, as well as delayed development of speech and language. Most of these mutations disrupt the function of the SHANK3 protein or lead to the production of an abnormally

short version of the protein. It is unclear how changes in the *SHANK3* gene are related to the risk of developing autism. Researchers suspect that a disruption in nerve cell communication contributes to the development of autism.

## Chromosomal Location

Cytogenetic Location: 22q13.33, which is the long (q) arm of chromosome 22 at position 13.33

Molecular Location: base pairs 50,674,642 to 50,733,212 on chromosome 22 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

## Other Names for This Gene

- proline-rich synapse-associated protein 2
- ProSAP2
- SPANK-2

## Additional Information & Resources

### Educational Resources

- Neuroscience (second edition, 2001): Synaptic Transmission  
<https://www.ncbi.nlm.nih.gov/books/NBK11001/>

### GeneReviews

- Phelan-McDermid Syndrome  
<https://www.ncbi.nlm.nih.gov/books/NBK1198>

### Scientific Articles on PubMed

- PubMed  
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SHANK3%5BTIAB%5D%29+OR+%28PROSAP2%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+2520+days%22%5Bdp%5D>

## OMIM

- SH3 AND MULTIPLE ANKYRIN REPEAT DOMAINS 3  
<http://omim.org/entry/606230>

## Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology  
[http://atlasgeneticsoncology.org/Genes/GC\\_SHANK3.html](http://atlasgeneticsoncology.org/Genes/GC_SHANK3.html)
- ClinVar  
<https://www.ncbi.nlm.nih.gov/clinvar?term=SHANK3%5Bgene%5D>
- HGNC Gene Family: Ankyrin repeat domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/403>
- HGNC Gene Family: PDZ domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/1220>
- HGNC Gene Family: Sterile alpha motif domain containing  
<http://www.genenames.org/cgi-bin/genefamilies/set/760>
- HGNC Gene Symbol Report  
[http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?q=data/hgnc\\_data.php&hgnc\\_id=14294](http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=14294)
- NCBI Gene  
<https://www.ncbi.nlm.nih.gov/gene/85358>
- UniProt  
<http://www.uniprot.org/uniprot/Q9BYB0>

## **Sources for This Summary**

- Boeckers TM, Bockmann J, Kreutz MR, Gundelfinger ED. ProSAP/Shank proteins - a family of higher order organizing molecules of the postsynaptic density with an emerging role in human neurological disease. J Neurochem. 2002 Jun;81(5):903-10. Review.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12065602>
- Bonaglia MC, Giorda R, Mani E, Aceti G, Anderlid BM, Baroncini A, Pramparo T, Zuffardi O. Identification of a recurrent breakpoint within the SHANK3 gene in the 22q13.3 deletion syndrome. J Med Genet. 2006 Oct;43(10):822-8. Epub 2005 Nov 11.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/16284256>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2563164/>
- Durand CM, Betancur C, Boeckers TM, Bockmann J, Chaste P, Fauchereau F, Nygren G, Rastam M, Gillberg IC, Anckarsäter H, Sponheim E, Goubran-Botros H, Delorme R, Chabane N, Mouren-Simeoni MC, de Mas P, Bieth E, Rogé B, Héron D, Burglen L, Gillberg C, Leboyer M, Bourgeron T. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. Nat Genet. 2007 Jan;39(1):25-7. Epub 2006 Dec 17.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17173049>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2082049/>

- Gauthier J, Spiegelman D, Piton A, Lafrenière RG, Laurent S, St-Onge J, Lapointe L, Hamdan FF, Cossette P, Motttron L, Fombonne E, Joobor R, Marineau C, Drapeau P, Rouleau GA. Novel de novo SHANK3 mutation in autistic patients. *Am J Med Genet B Neuropsychiatr Genet*. 2009 Apr 5; 150B(3):421-4. doi: 10.1002/ajmg.b.30822.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/18615476>
- Moessner R, Marshall CR, Sutcliffe JS, Skaug J, Pinto D, Vincent J, Zwaigenbaum L, Fernandez B, Roberts W, Szatmari P, Scherer SW. Contribution of SHANK3 mutations to autism spectrum disorder. *Am J Hum Genet*. 2007 Dec;81(6):1289-97. Epub 2007 Oct 16.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/17999366>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2276348/>
- OMIM: SH3 AND MULTIPLE ANKYRIN REPEAT DOMAINS 3  
<http://omim.org/entry/606230>
- Wilson HL, Wong AC, Shaw SR, Tse WY, Stapleton GA, Phelan MC, Hu S, Marshall J, McDermid HE. Molecular characterisation of the 22q13 deletion syndrome supports the role of haploinsufficiency of SHANK3/PROSAP2 in the major neurological symptoms. *J Med Genet*. 2003 Aug;40(8):575-84.  
*Citation on PubMed:* <https://www.ncbi.nlm.nih.gov/pubmed/12920066>  
*Free article on PubMed Central:* <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1735560/>

---

Reprinted from Genetics Home Reference:  
<https://ghr.nlm.nih.gov/gene/SHANK3>

Reviewed: September 2009  
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications  
U.S. National Library of Medicine  
National Institutes of Health  
Department of Health & Human Services